

Report No. : TCT171220C019 Date : Jan. 03, 2018 Page No.: 1 of 6 Applicant: Shenzhen LTQ VAPOR Electronics Co., Ltd Address: 5F/L, Building 1, FuHong Industrial Park, Fuyong Town, Baoan District, Shenzhen, China The following sample was submitted and identified by/on behalf of the client as: DEMON KILLER TINY KIT Sample Name: Model No.: DEMON KILLER TINY KIT Tank: SS316+PEI Voltage level in testing: Voltage/Wattage of tested sample is un-adjustable Adjustable air inlet or not: Yes Trade Mark: DEMON KILLER Sample Received Date: 2017.12.25 **Testing Period:** 2017.12.25-2018.01.03 As specified by client, to determine the Carbonyl Compounds content(s) Test Requested: 1. in aerosol generated by the submitted sample. 2. As specified by client, to determine the Metals content(s) in aerosol generated by the submitted sample. 3. As specified by client, to determine Nicotine consistency in aerosol generated by the submitted sample. Please refer to the following page(s). Test Method: Please refer to the following page(s). Test Result(s): Checked by Signed for and on behalf of TCT Chris Zhong Kim Zhang **Technical Manager** Shenzhen TCT Testing Technology Co., Ltd.



Report No. : TCT171220C019

Date : Jan. 03, 2018

Page No.: 2 of 6

Test Results:

Test Condition for test items except Nicotine consistency test:

With reference to the CORESTA RECOMMENDED METHOD Nº 81 method parameter and Afnor standardization XP D90-300-3, a smoke machine was used to collect the vapor.

Puff Duration	3.0s±0.1s
Puff Volume	55mL±0.3mL
Puff Frequency	30s±0.5s
Puff of Each Group	20
Group Interval Time	300s±120s
Maximum Flow	18.5mL/s±1.0mL/s
Pressure Drop	< 50hPa
Group	5
Total Number of Puff	100
Total Duration of Vaporization	300s

The temperature and relative humidity of the test atmosphere during machine preparation and testing shall be kept within the following limits: temperature $\pm 2^{\circ}$ C, relative humidity $\pm 5^{\circ}$

Sample Description:

1. DEMON KILLER TINY KIT



Report No. : TCT171220C019

Date : Jan. 03, 2018

Page No.: 3 of 6

1. Carbonyl Compounds Content(s)

Method: Using volumes based on the desired dilution, a measured volume of sample was combined with a volume of DNPH solution and vortexed. After sitting for 20 minutes at ambient temperature, the sample was then quenched with a sufficient amount of pyridine. An aliquot was then analyzed using the Agilent Model 1200, High Performance Liquid Chromatograph equipped with an Ultraviolet (UV) Detector operating at 365 nm.

Test Item	CAS No.	CAS No. Unit		LOQ	Content(s) 1	
Formaldehyde	50-00-0	ug/100puffs	0.667	2	14.5	
Acetaldehyde	75-07-0	ug/100puffs	0.667	2	ND	
Acrolein	107-02-8	ug/100puffs	0.667	2	ND	

- Note: ug = Microgram
 - ND = Not Detected (lower than MDL)
 - MDL = Method Detection Limit
 - LOQ = Limit of Quantitation
 - E-Liquid Used: E-liquid B (AFNOR XP D90-300-3)



Report No. : TCT171220C019

Date : Jan. 03, 2018

Page No.: 4 of 6

2. Metals Content(s)

Method: The vapor was passed through a dry-ice cooled impinger containing glass packing beads and quartz wool. After smoking the impinger was extracted with 5% nitric acid and filtered through quartz wool. An aliquot of the resulting solution was submitted for analysis by ICP-OES.

Test Item	CAS No.	Unit	MDL	LOQ	Content(s)
Aluminium(Al)	7429-90-5	ug/100puffs	0.025	0.25	ND
Chromium(Cr)	7440-47-3	ug/100puffs	0.005	0.05	ND
Iron(Fe)	7439-89-6	ug/100puffs	0.005	0.05	ND
Nickel(Ni)	7440-02-0	ug/100puffs	0.025	0.25	ND
Tin(Sn)	7440-31-5	ug/100puffs	0.25	2.5	ND

Note: - ug = Microgram

- ND = Not Detected (lower than MDL)
- MDL = Method Detection Limit
- LOQ = Limit of Quantitation
- E-Liquid Used: E-liquid B (AFNOR XP D90-300-3)



Report No. : TCT171220C019

Date : Jan. 03, 2018

Page No.: 5 of 6

3. Nicotine Consistency Test

Test Condition: With reference to the CORESTA RECOMMENDED METHOD Nº 81 method parameter and Afnor standardization XP D90-300-3, a smoke machine was used to collect the vapor.

Puff Duration	3.0s±0.1s			
Puff Volume	55mL±0.3mL			
Puff of Each Group	20			
Maximum Flow	18.5mL/s±1.0mL∕s			
Pressure Drop	< 50hPa			

The temperature and relative humidity of the test atmosphere during machine preparation and testing shall be kept within the following limits: temperature $\pm 2^{\circ}$ C, relative humidity $\pm 5\%$

Method: A reference liquid was prepared. A pharmaceutical nicotine inhaler was used as a comparator. Products were attached to a smoke machine, and the aerosol was collected in Cambridge filter pads. After trapping and solvent extraction, solution was analyzed by GC-MS and nicotine was dosed by comparing the areas obtained on the MS detector with those of standard solutions prepared in the laboratory under concentration conditions surrounding those of the samples.

Sample	Nicotine(CAS No.:54-11-5) Contents(mg / 20 Puffs)					Total(mg/	
Description	Group 1*	Group 2	Group 3*	Group 4	Group 5*	AVG	100puffs)
DEMON KILLER TINY KIT	0.728	0.707	0.683	0.702	0.706	0.705	3.53
Deviation(%)	3.2	-	3.1	-	0.1	-	-

Note: - mg = milligram

- ND = Not Detected (lower than MDL)
- MDL = Method Detection Limit = 0.01 mg / 20 Puffs
- LOQ = Limit of Quantitation = 0.1 mg / 20 Puffs
- 1group = 20 puffs
- * Values used for determination of consistency of nicotine emission
- E-Liquid Used: E-liquid A (AFNOR XP D90-300-3)
- Under the conditions of the test and with reference to AFNOR XP D90-300-3, the electronic cigarette delivers a dose of nicotine at consistent levels.



Report No. : TCT171220C019

Date : Jan. 03, 2018

Page No.: 6 of 6

Photo(s) of the sample(s)



Remark: This report is considered invalidated without the Special Seal for Inspection of the TCT. This report shall not be altered, increased or deleted. The results shown in this test report refer only to the sample(s) tested. Without written approval of TCT, this test report shall not be copied except in full and published as advertisement.